

Global implications of 1.5 °C and 2 °C warmer worlds on extreme river flows

November 7th, 2018



Photo: Video footage in a museum in Florence, Italy, shows the 5 m high flood marks on the wall in this museum after the 1966 river flood. Extreme river flows will occur more frequent in most parts of the world.

The Paris targets agreed upon in 2015 aim to limit global warming to $\frac{1}{2}$ well below 2 °C and to pursue e®orts to limit the temperature increase to 1.5 °C above pre-industrial levelsI. The impacts of these global warming scenarios on global peak river 2 ows was studied for peak 2 ows with a 100 year return period.

In most world regions peak river flows will occur more frequent

According to this study, in most regions of the world peak river 2ows that currently occur once every 100 years occur more frequent at 1.5 °C global warming, and this frequency further increases at 2.0 °C warming. These regions include South America (Amazon catchment), central Africa, central-western Europe, the south of the US (Mississippi river area), central Asia, and Siberian catchments. In these regions, the frequency of the current 1in-100 year 2ow typically shifts to once in 70-90 years at 1.5 °C, and once in 50 years at 2.0 °C global warming. The current 1-in-100 year 2ow becomes even more common in most of central and eastern China, the southern part of the Amazon catchment, the Blue Nile, and in northern India: once in 50-60 years at 1.5 °C warming, and once in 25-35 years at 2 °C warming. Frequency changes are small for North America and eastern Europe.

In some world regions peak river flows will occur less frequent

Some world regions see a decrease in the frequency of high 2ows between the present day and 1.5 °C or 2.0 °C warming. These regions are located in the northern part of South America, areas of the western coast of the United States (Colorado River) and Canada, South Africa (Orange River), Scandinavia, and in some parts of eastern Europe. In most of these regions, according to this study, the current 1-in-100 year 2ow decreases in frequency to approximately 1-in-150 years, with little further decrease at 2.0 °C.

Why peak river flow frequency will change

The authors of this study relate these changes to a decreasing role of snowmelt in the generation of high 20ws as snow accumulation reduces, and to a more dominant role of extreme precipitation. The latter results in a progressive increase in the frequency of high 20ws in large parts of the world, despite a reduction in the number of rain days. This suggests an intensilcation of hydrological extremes in these regions. In fact, more frequent peak river 20ws and more frequent droughts can go hand in hand: in some world regions both an increase in high 20w frequencies and a decrease in mean river 20w accompanied by major drought hazards is projected for 2.0 °C global warming.

Source: Paltan et al. (2018). Environmental Research Letters 13.