

# In a changing climate, the 100year flood event is a misleading concept

July 26th, 2021



In the media, river stoods are often characterized by a return period, a long return period being an indication that the stood was exceptionally extreme. A long return period means a low probability of that particular stood in any year. The stood with a return period of 100 years is mostly used as a benchmark to characterize the extreme nature of any stood.

### 100-year flood event: misleading concept

In a changing climate, concepts like these '100-year stood events' can be misleading, however. The discharge linked to a 100-year return period is based on a historical record. But the character of rivers is changing, not only in response to the changing climate but also as a result of changes in land use, such as urbanization and agriculture, and other anthropogenic instuences on river regimes. Data of the past may no longer fit the river's discharge regime of today.

Basically, this means that statements on a river sbod's return period to indicate its extreme character are of little value. Unless these statements are based on a trend analysis of sbod levels over a sufficiently long period.

#### Global assessment

The latter has been carried out on a global scale. Changes have been examined in return periods of observed extreme peak river sows globally. According to the authors, they provide the first global picture of changes in the return periods of the 20-, 50-, and 100-year river soods. Their reference point is 1970. The analysis is based on more than 10,000 river gauge records. At each site, the authors estimated the magnitude of the 20-, 50-, and 100-year sood in the 1970s, and the changing return period of these 1970s soods in the current situation.

#### Western Europe: reduction return period

With the most recent stoodings in Western Europe in mind, most people will think that the return period of extreme stoods has reduced, and their probability of occurrence has increased. Indeed, in temperate climate zones including Western Europe, return periods have decreased for the 20- and 50-year stoods. On average, in temperate climates globally, stoods that had a return period of 20 and 50 years in 1970, now have return periods of 8 and 21 years, respectively.

According to the analysis, the return period of a 100-year shoot has increased. One should be aware, however, that the number of sites used to evaluate these trends is much smaller for the 100-year shoots than for the 20-year shoots. After all, the numbers of sites with a sufficiently long record to carry out these analyses are much smaller for increasingly rare extremes. The conclusions on the 100-year-shoots are, therefore, of little value.

#### The global picture: increase return period

Remarkably, the overall global impression of a changing return period is not a reduction, but an increase of these shod return periods: The 20-, 50-, and 100-year shods of the 1970s have become, on average, 41-, 152-, and 358-year shods under present-day conditions. With the exception of temperate zones, this increase of return period was found for all other climate zones (cold, arid, tropical, polar) and includes eastern Europe.

This indicates that, on a global scale, the frequency of occurrence of these shool levels has decreased. According to the authors, this may be due to a combination of climatic changes, land cover changes such as urbanization, agriculture, and other anthropogenic instuences on river regimes.

#### An indication, no precise estimates

The authors stress that these 'new' return periods should be seen as indicative of the general direction of change rather than precise estimates. After all, the results only cover the selected sites with sufficiently long historic shoot records.

## Climate change versus catchment alterations

Previously, we reported on a global study on 20-, 50-, and 100-year stoods where the effects of human catchment alterations, including urbanization, land-use change, diversions, abstractions and reservoir regulation, were separated from those of climate change. In this way, only the impact of climate change on river stoods could be studied. This study showed no compelling evidence for consistent changes over time in the return period of these stoods.

Source: Slater et al., 2021. Geophysical Research Letters 48.