

Accelerated dryland expansion under climate change

March 5th, 2016



The global area of dryland is increasing rapidly. This was shown from data over the period 1948–2005, and seems to proceed towards the end of this century, according to projections of a large number of climate models.

The aridity of a region is generally measured by the aridity index (AI), which is the ratio of total annual precipitation to potential evapotranspiration. Drylands are defined as regions with $AI < 0.65$.

Dryland expansion rate will increase in future decades: according to projections of 20 global climate models, global dryland area may increase by 11% and 23% by the end of the twenty-first century under a moderate and high-end scenario of climate change (the so-called RCP4.5 and RCP8.5 scenario), respectively, relative to the reference period of 1961–1990. As a result dryland will cover the global land surface by 50% - 56% in 2100. Under the high-end

scenario of climate change, 78% of dryland expansion may occur in developing countries. The increasing aridity, enhanced warming and rapidly growing human population will exacerbate the risk of land degradation and desertification in these countries in the near future.

Drylands are divided into subtypes of hyper-arid ($AI < 0.05$), arid ($0.05 \leq AI < 0.2$), semiarid ($0.2 \leq AI < 0.5$) and dry subhumid ($0.5 \leq AI < 0.65$) regions (Middleton and Thomas, 1997). During the period 1948–2005 the observed global land area increases in hyper-arid, arid, semiarid and subhumid land types are 0.62%, 1.16%, 2.32% and 3.32%. Under the moderate scenario of climate change the dryland subtypes of dry subhumid, semiarid, arid and hyper-arid comprise 8.9%, 19.0%, 14.4% and 8.4% of the global land area in 2100. Under the high-end scenario of climate change these subtypes of dry subhumid, semiarid, arid and hyper-arid comprise 8.3%, 20.3%, 14.9% and 12.6% of the global land area.

Strict management and rational utilization of water resources, along with the restoration of soils and vegetation to reduce ecosystem vulnerability on a global scale, are urgently needed to develop a global action plan to prevent future desertification and eradicate the present global desertification problems.

Source: Huang et al., 2016. *Nature Climate Change* 6: 166-172.

Photo: Ingo Vogelmann (www.flickr.com)