

Vulnerability European cities to extreme events is increasing

March 6th, 2017



Already in the next decades highly populated urban areas in Central Europe will experience significantly more hot days, tropical nights, and extreme precipitation events. This was shown for an area covering the east of France, Germany, Austria, Czech Republic, Slovenia, and the north of Italy. The study was based on high-resolution climate simulations and a scenario of moderate climate change for the period 2021-2050, compared to a reference period 1971-2000.

According to this study Central Europe will show an average increase of the number of heat waves (17%), as well as the number of single hot days (12%), tropical nights (30 %) and heavy precipitation events (40%), and a decrease of the number of frost days (30%). For both hot and frost days absolute decrease is largest in the northeast whereas relative changes are largest in the northwest.

The results show that the number of hot days (daily maximum temperatures $> 30^{\circ}\text{C}$) will increase particularly in the western part of central Europe, affecting cities like Paris, Stuttgart and Hamburg. At some locations, the number of hot days is to increase by over 70%. The

number of tropical nights (daily minimum temperature $> 20^{\circ}\text{C}$) shows a similar pattern since 'hot days' regions cannot significantly cool down and high temperatures remain throughout the night. The relative increase of the number of tropical nights shows a clear trend from west to east, ranging from roughly 70% for Hamburg and 40% for Paris, down to less than 10% in the vicinity of Vienna. Also the increase in the number of heat waves is higher in the western part (with cities like Paris and Zurich) than in the eastern part.

In general, a decrease of the number of frost days is projected, by up to 45% at the coastline of the Atlantic Ocean. Remarkably, there is no clear projected change in the number of frost days in the alpine area.

The increase of the number of days that buildings have to be air-conditioned in the summer dominates over the decrease that buildings need to be heated in the winter.

In Europe heavy precipitation events, with daily accumulated precipitation > 20 mm, are most likely in the summer (June, July and August). For Central and Northwestern Europe the relative increase in the number of days with heavy precipitation in these months is largest in northern Germany, Belgium and the Netherlands ($> 30\%$).

Source: Fallmann et al., 2017. Theoretical and Applied Climatology 127: 667-683

Photo: Gary Knight (www.flickr.com)